

Structure Function Scaling in Compressible Super-Alfvénic MHD Turbulence

Paolo Padoan¹, Raul Jimenez², Åke Nordlund³ and Stanislav Boldyrev⁴

¹*Physics Department, University of California, San Diego, La Jolla, CA 92093*

²*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104*

³*Copenhagen Astronomical Observatory and Theoretical Astrophysics Center, DK-2100, Copenhagen, Denmark*

⁴*Department of Astronomy and Astrophysics, University of Chicago, 5640 S. Ellis Avenue, Chicago, IL 60637*

(Dated: February 2, 2008)

Supersonic turbulent flows of magnetized gas are believed to play an important role in the dynamics of star-forming clouds in galaxies. Understanding statistical properties of such flows is crucial for developing a theory of star formation. In this letter we propose a unified approach for obtaining the velocity scaling in compressible and super-Alfvénic turbulence, valid for arbitrary sonic Mach number, M_S . We demonstrate with numerical simulations that the scaling can be described with the She-Lévêque formalism, where only one parameter, interpreted as the Hausdorff dimension of the most intense dissipative structures, needs to be varied as a function of M_S . Our results thus provide a method for obtaining the velocity scaling in interstellar clouds once their Mach numbers have been inferred from observations.

PACS numbers: 98.38.Am, 95.30.Qd, 98.38.Dq, 95.30.Lz

1. Introduction. Star formation occurs in our galaxy primarily inside cold interstellar clouds. Observations indicate that gas motion inside these clouds is highly supersonic in the broad range of scales $0.1 - 1000$ pc. At these scales the clouds lack any characteristic structure and their scaling properties are to some extent universal. This universality is the signature of the supersonic and super-Alfvénic magneto-hydrodynamic (MHD) turbulence that governs the dynamics of the clouds [1, 2, 3, 4, 5, 6, 7].

The properties of turbulent MHD flows are controlled by the rms sonic Mach number, M_S , (ratio of rms flow velocity and sound velocity) and the rms Alfvénic Mach number, M_A , (ratio of rms flow velocity and Alfvén velocity). In interstellar clouds these numbers are large, $M_S \sim 1 - 30$, $M_A > 1$. The corresponding supersonic motion produces the observed strong density enhancements [8], some of which are Jeans-unstable and undergo gravitational collapse. This initial ‘turbulent’ stage of density fragmentation should be described statistically, in terms of velocity and density probability distribution functions or correlators; its properties are inherently related to the statistical properties of compressible MHD turbulence [9].

Early studies suggesting the importance of astrophysical supersonic turbulence were undertaken by von Weizsäcker [10] and later by Larson [11], who discovered that the spectrum of the velocity field in molecular clouds was steeper than the spectrum predicted by the standard Kolmogorov model. The discrepancy was attributed to the strong intermittency of supersonic turbulence. However, the analytical understanding of such turbulence remained rather poor.

Advances in computational and observational techniques have recently allowed measurements of power spectra and higher order correlators of velocity and density fields, reviving the interest in supersonic turbulence. Modern observations generate surveys covering a range

of scales comparable or even exceeding the resolution of numerical simulations. The interstellar medium can now be viewed as a laboratory for investigating supersonic motion unachievable with Earth-based experiments.

The turbulent field can be characterized by the scaling of its structure functions [see below]. The intermittency is defined as the departure of this scaling from the Kolmogorov one. So far, only the limiting cases of $M_S \gg 1$ and $M_S = 0$ have been investigated. An analytical description of supersonic turbulence, $M_S \gg 1$, has been proposed in [12] and confirmed numerically in [13, 14, 15]. It is based on the observation that the most intense dissipative structures in supersonic flows are two-dimensional shocks. This approach makes use of the model by She and Lévêque [16] for incompressible turbulence, $M_S = 0$, where the dissipative structures are one-dimensional vortex filaments.

In this letter we report the results of a series of numerical MHD simulations where we traced the change of the velocity structure functions over the broad range of Mach numbers $0.3 \leq M_S \leq 10$. We discovered that their scalings can be described by the general model where only one parameter, the dimension of the most intense dissipative structures, needs to be changed with M_S . The scalings obtained in [12, 16] naturally appear as the limiting cases of this unifying model. Our results suggest that the velocity scaling within interstellar clouds can be inferred from measurements of their Mach number. Observationally, this is a much easier task than measuring the structure function directly. We present the main results here; the detailed discussion will appear elsewhere.

2. Velocity Structure Function Scaling. The velocity structure functions of order p are defined as:

$$S_p(\ell) = \langle |u(x + \ell) - u(x)|^p \rangle \propto \ell^{\zeta(p)} \quad (1)$$

where the velocity component u is parallel (longitudinal structure function) or perpendicular (transversal struc-




FIG. 1: Compensated velocity structure functions of order $p = 1$ to $p = 10$ (bottom to top) from the $M_S=9.5$ experiment. To avoid confusion, the structure functions of order $p = 1$ and $p = 2$ have been multiplied by 0.2 and 0.7 respectively and only the upper part of error bars larger than 100% is shown.

ture function) to the vector ℓ and the spatial average is over all values of the position x .

The scaling of velocity structure functions in incompressible turbulence is best described by the She–Lévêque formula [16, 17]:

$$\zeta(p)/\zeta(3) = p/9 + 2 \left[1 - (2/3)^{p/3} \right], \quad (2)$$

This is a specific case of the more general formula

$$\zeta(p)/\zeta(3) = \gamma p + C(1 - \beta^p) \quad (3)$$

with the normalization condition,

$$C = (1 - 3\gamma)/(1 - \beta^3). \quad (4)$$

Two basic assumptions are used to derive (3), as discussed in [18]. The first assumption is the existence of a symmetric relation between the intensity of fluctuations $F_p(\ell)$ and $F_{p+1}(\ell)$ with respect to a translation in p :

$$F_{p+1}(\ell) = A_p F_p(\ell)^\beta F_{(\infty)}(\ell)^{1-\beta}, \quad (5)$$

where the p th order intensity of fluctuations is defined as $F_p(\ell) = S_{p+1}(\ell)/S_p(\ell)$, and A_p are constants independent of ℓ (generally found to be independent of p as well). The functions $F_p(\ell)$ are associated with higher intensity fluctuations with increasing values of p . The parameter β is a measure of intermittency. In Kolmogorov’s 1941 model [19], $\beta \rightarrow 1$ and $\zeta(p) = p/3$. This is the limit of no intermittency. Lower β corresponds to higher degree of intermittency, or to the increasing role played by strong persistent fluctuations (structures).




FIG. 2: Structure function exponents normalized to the values predicted by [12] for supersonic turbulence (dashed–dotted line). Values from five experiments are shown, from $M_S=9.5$ (bottom) to $M_S=0.4$ (top). The $M_S=0.8$ case is not shown as it almost overlaps with the $M_S=0.4$ experiment.

The second assumption is

$$F_\infty \sim S_3^\gamma. \quad (6)$$

The value of the parameter γ is related to very high order moments and is therefore difficult to constrain with experiments. Kolmogorov’s turbulence corresponds to $\gamma = 1/3$.

The parameter C has been interpreted as the Hausdorff codimension of the support of the most singular dissipative structures [17]. In incompressible turbulence the most dissipative structures are along vortex filaments with Hausdorff dimension $D = 1$ and so $C = 2$. Boldyrev [12] has proposed to extend the scaling (3) to supersonic turbulence, with the assumption that the Hausdorff dimension of the support of the most singular dissipative structures is $D = 2$ ($C = 1$), because dissipation of supersonic turbulence occurs mainly in sheet-like shocks. He used $\gamma = 1/9$ as in incompressible turbulence, which implies $\beta^3 = 1/3$, that is a higher degree of intermittency than in incompressible turbulence. With these parameters one obtains the velocity scaling proposed for supersonic turbulence in [12]:

$$\zeta(p)/\zeta(3) = p/9 + 1 - (1/3)^{p/3}. \quad (7)$$

This velocity scaling has been found to provide a very accurate prediction for numerical simulations of highly supersonic and super-Alfvénic turbulence [13, 14]. The same scaling has also been proposed for incompressible MHD turbulence [20], where dissipation occurs mainly in two dimensional current sheets.




FIG. 3: “ β test” plot for the models with $M_S=0.4$ and $M_S=9.5$. The $M_S=0.4$ model is shifted upward by 0.2 y-units. The solid lines are least square fits with slope β .

In the present work we obtain the structure function scaling in simulations of compressible super-Alfvénic MHD turbulence as a function of the rms sonic Mach number of the flow, M_S , for constant value of the rms Alfvénic Mach number, $M_A \gg 1$ (the limiting case of strong magnetic field was investigated analytically and numerically in [21, 22, 23]). We use an isothermal equation of state and vary the value of M_S in different experiments by varying the thermal energy. As a result, the initial value of $M_A=10$ remains unchanged from run to run. In our numerical method the total magnetic flux is conserved. However, the magnetic energy is amplified.

The initial value of the ratio of average magnetic and dynamic pressures is $\langle P_m \rangle_{\text{in}} / \langle P_d \rangle_{\text{in}} = 0.005$ for all runs.

The value of the same ratio, averaged over the time interval for which the structure functions are computed, is given in Table 1. Because the average dynamic pressure is always in excess of the average magnetic pressure, $\langle P_m \rangle / \langle P_d \rangle \ll 1$, the dissipation is expected to occur primarily in vortex filaments (roughly corresponding to magnetic flux tubes) in subsonic runs and in shocks in supersonic runs. In the next section we argue that when we increase the value of M_S , the Hausdorff dimension of the support of the most dissipative structures grows continuously from $D \approx 1$ to $D \approx 2$.

3. Results. We have solved the three dimensional compressible MHD equations, for an isothermal gas with initially uniform density and magnetic fields, in a staggered mesh with 250^3 computational cells and periodic boundary conditions. Turbulence is set up as an initial large scale random and solenoidal velocity field (generated in Fourier space with power only in the range of wavenumber $1 \leq K \leq 2$) and maintained with an exter-




FIG. 4: Hausdorff dimension of the support of the most dissipative structures versus rms sonic Mach number.

nal large scale random and solenoidal force, correlated at the largest scale turn-over time. Details about the numerical method are given in [3].

We have run six experiments with different values of M_S , in the range $0.4 \leq M_S \leq 9.5$ (Table I), for approximately 10 dynamical times. The transverse structure functions of velocity are computed for the last four dynamical times of each experiment, up to the order $p = 10$. Their power law slope, $\zeta(p)$, is obtained from a least square fit in the range $2 \leq \ell \leq 96$. The standard deviation of the slope is used as an estimate of the uncertainty of $\zeta(p)$.

Figure 1 shows the structure functions plotted versus the third order one for the $M_S=9.5$ experiment. The structure functions are compensated by their least square fit slope, $\zeta(p)$. We study the structure functions relative to the third order to exploit the concept of extended self-similarity [17, 24], that is the property of the relative scaling to extend up to a range of scales affected by dissipation (numerical dissipation in this case).

The exponents $\zeta(p)$, normalized to the third order one, are plotted in Figure 2, divided by their value predicted in [12] and given by (7). The $M_S=0.4$ and $M_S=0.8$ models follow the She-L  v  que scaling of incompressible turbulence. As the value of M_S increases, the velocity scaling becomes gradually more intermittent. At $M_S=9.5$ the scaling is very close to the analytical prediction for supersonic turbulence [12], as previously shown [13, 14].

The validity of the assumption of hierarchical symmetry expressed by (5), can be tested with a log-log plot of F_{p+1}/F_2 versus F_p/F_1 , obtained by varying the argument ℓ in (5). If the plot is a straight line, the parameters A_p are constants independent of ℓ and the assumed hierarchical symmetry expressed by (5) is satisfied. This

$\langle M_S \rangle$	0.39	0.77	1.5	2.6	4.8	9.5
$\langle P_m \rangle_{\text{in}} / \langle P_d \rangle_{\text{in}}$	0.005	0.005	0.005	0.005	0.005	0.005
$\langle P_m \rangle / \langle P_d \rangle$	0.15	0.14	0.09	0.11	0.09	0.08
β	0.87	0.87	0.84	0.79	0.73	0.69
D	1.0	1.0	1.34	1.70	1.90	2.01

TABLE I: Values of the rms sonic Mach number, average magnetic to dynamic pressure ratio at initial time and averaged over the time interval for which the structure functions are computed, β parameter from fitting $\zeta(p)/\zeta(3)$ and Hausdorff dimension of the support of the most dissipative structures.

is called the “ β test” in [18]. Our numerical simulations pass the β test for all values of M_S . In Figure 3 we show the two extreme cases, $M_S=0.4$ and $M_S=9.5$. We have used values of $F_p(\ell)$ for all values of ℓ used in the least square fit estimate of the exponents $\zeta(p)$. In all the runs we find that the constants A_p in (5) are independent not only of ℓ , but also of p , as in previous numerical and laboratory studies [18, 25, 26]. The plot in Figure 3 is in fact obtained varying both ℓ and p .

The slope of the plot in Figure 3 provides an estimate of β . For the models with M_S from 0.4 to 9.5 we obtain $\beta = 0.82, 0.81, 0.75, 0.74, 0.77$ and 0.77 . We instead find no evidence of a systematic variation in the parameter γ with M_S and our data is consistent with $\gamma = 1/9$.

The β test gives support to the basic assumption of the She–L  v  que formalism, but does not necessarily yield the value of β providing the best fit to the structure function scaling $\zeta(p)/\zeta(3)$. Values of β computed from a best fit to $\zeta(p)/\zeta(3)$ from our numerical experiments are slightly different from the β test values (see Table I). We use these best-fit β values to compute the parameter D .

From the definition of the parameter C in (4), the values of β estimated above and $\gamma = 1/9$ we can compute $D = 3 - C$ as a function of M_S . The values of β and D are given in Table I. The value of D grows with M_S , from $D \approx 1.0$ in the subsonic runs to $D \approx 2.0$ in the most supersonic runs.

4. Conclusions. We propose a unifying description of the velocity structure function scaling of compressible and super-Alfv  nic MHD turbulence for an arbitrary value of the rms sonic Mach number of the flow, M_S . Only one model parameter, interpreted as the Hausdorff dimension of the support of the most dissipative structures in the She–L  v  que formalism, needs to be varied as a function of M_S . Besides its general importance, our result implies that the velocity structure function scaling in interstellar clouds can be predicted because the value of M_S in the clouds is easily estimated from observations. This may help improve our understanding of the process of cloud fragmentation that leads to the birth of stars.

RJ was supported by NSF grant AST-0206031.   N was supported by a grant from the Danish Natural Science Research Council, and in part by the Danish National Research Foundation, through its establishment of the Theoretical Astrophysics Center. SB was supported by the NSF Center for magnetic self-organization in astrophysical and laboratory plasmas at the University of Chicago.

-
- [1] P. Padoan, J. Bally, Y. Billawala, M. Juvela, and   . Nordlund, *ApJ* **525**, 318 (1999).
 - [2] J. Ballesteros-Paredes, L. Hartmann, and E. V  zquez-Semadeni, *ApJ* **527**, 285 (1999).
 - [3] P. Padoan and   . Nordlund, *ApJ* **526**, 279 (1999).
 - [4] R. S. Klessen, F. Heitsch, and M. Mac Low, *ApJ* **535**, 887 (2000).
 - [5] E. C. Ostriker, J. M. Stone, and C. F. Gammie, *ApJ* **546**, 980 (2001).
 - [6] P. Padoan, L. Cambresy, and W. Langer, *ApJ*, in press (astro-ph/0208217) (2002).
 - [7] P. Padoan, S. Boldyrev, W. Langer, and   . Nordlund, *ApJ*, in press (astr-ph/0207568) (2002).
 - [8] P. Padoan, M. Juvela, A. A. Goodman, and   . Nordlund, *ApJ* **553**, 227 (2001).
 - [9] P. Padoan and   . Nordlund, *Astrophys. J.* **576**, 870 (2002).
 - [10] C. F. von Weizs  cker, *ApJ* **114**, 165 (1951).
 - [11] R. B. Larson, *MNRAS* **194**, 809 (1981).
 - [12] S. Boldyrev, *ApJ* **569**, 841 (2002).
 - [13] S. Boldyrev,   . Nordlund, and P. Padoan, *Astrophys. J.* **573**, 678 (2002).
 - [14] S. Boldyrev,   . Nordlund, and P. Padoan, *Physical Review Letters* **89**, 031102 (2002).
 - [15] A. G. Kritsuk and M. L. Norman, *astro-ph/0310697* (2003).
 - [16] Z.-S. She and E. L  v  que, *Phys. Rev. Lett.* **72**, 336 (1994).
 - [17] B. Dubrulle, *Phys. Rev. Lett.* **73**, 959 (1994).
 - [18] Z.-S. She, K. Ren, G. S. Lewis, and H. L. Swinney, *Phys. Rev. E* **64**, 016308 (2001).
 - [19] A. N. Kolmogorov, *Dokl. Akad. Nauk. SSSR* **30**, 301 (1941).
 - [20] W.-C. M  ller and D. Biskamp, *Phys. Rev. Lett.* **84**, 475 (2000).
 - [21] Y. Lithwick and P. Goldreich, *Astrophys. J.* **562**, 279 (2001).
 - [22] J. Cho, A. Lazarian, and E. T. Vishniac, *Astrophys. J.* **564**, 291 (2002).
 - [23] J. Cho and A. Lazarian, *Physical Review Letters* **88**, 245001 (2002).
 - [24] R. Benzi, S. Ciliberto, R. Tripiccion, C. Baudet, F. Massaioli, and S. Succi, *Phys. Rev. E* **48**, 29 (1993).
 - [25] E. L  v  que and Z.-S. She, *Phys. Rev. E* **55**, 2789 (1997).
 - [26] G. R. Chavarr  a, C. Baudet, and S. Ciliberto, *Physical Review Letters* **74**, 1986 (1995).